

### GENETIC TESTING: KIDNEY DISORDERS

MP9598

Covered Service:	Yes
Prior Authorization Required:	No
Additional Information:	Use the current applicable CPT/HCPCS code(s). An appropriate diagnosis code must appear on the claim. Claims will deny in the absence of applicable diagnosis and procedure code(s) and/or if the criteria for coverage outlined below are not met. The following codes are included below for informational purposes only, and may be subject to change without notice. Inclusion or exclusion of a code does not constitute or imply member coverage or provider reimbursement.
Medicare Policy:	Prior authorization is dependent on the member's Medicare coverage. Prior authorization is not required for Dean Care Gold and Select when this service is provided by participating providers. If a member has Medicare primary and Dean Health Plan as secondary coverage, a prior authorization is required.
BadgerCare Plus Policy:	Dean Health Plan covers when BadgerCare Plus also covers the benefit.

Dean Health Plan Medical Policy:

# **OVERVIEW**

Inherited kidney disorders and inherited disorders that indirectly affect the kidneys can be more common, such as autosomal dominant polycystic kidney disease, or more rarely Lowe syndrome and Fabry disease. Identifying the genetic cause of an inherited kidney disorder can help direct treatment, inform family members, and contribute to the overall understanding of the genetic etiology of chronic kidney disease. More advanced next-generation sequencing, such as exome sequencing and comprehensive genetic testing panels, are emerging as a first-line diagnostic method for patients with chronic kidney disease.



# POLICY REFERENCE TABLE

The tests and associated laboratories and CPT codes contained within this document serve only as examples to help users navigate claims and corresponding coverage criteria; as such, they are not comprehensive and are not a guarantee of coverage or non-coverage. Please see the <u>Concert Genetics</u> <u>Platform</u> for a comprehensive list of registered tests.

<u>Coverage</u> <u>Criteria</u> <u>Sections</u>	Example Tests (Labs)	Common CPT Codes	Common ICD Codes	<u>Ref</u>		
Polycystic Kidney Disease						
<u>Targeted Variant</u> <u>Analysis</u>	Targeted Mutation Analysis for a Known Familial Variant	81403		8		
<u>Single gene or</u> <u>Multigene Panel</u>	Autosomal Dominant Polycystic Kidney Disease via the PKD1 Gene (PreventionGenetics, part of Exact Sciences)	81407, 81479	Q61, N18	1, 2		
	PKD2 Full Gene Sequencing and Deletion/Duplication (Invitae)	81406, 81479				
	Autosomal Recessive Polycystic Kidney Disease (ARPKD) via the PKHD1 Gene (PreventionGenetics, part of Exact Sciences)	81408, 81479				
	Autosomal Dominant Polycystic Kidney Disease (ADPKD) via the GANAB Gene (PreventionGenetics, part of Exact Sciences)	81479				
	Autosomal Dominant Polycystic Kidney Disease (ADPKD) via the DNAJB11 Gene (PreventionGenetics, part of Exact Sciences)					
	Hereditary Cystic Kidney Diseases Panel (PreventionGenetics, part of Exact Sciences)	81404, 81405, 81406, 81407, 81408, 81479				
	Polycystic Kidney Disease Panel (GeneDx)					
Comprehensive Kidney Disease Panels						
	RenaSight (Natera)			3		



<u>Coverage</u> <u>Criteria</u> <u>Sections</u>	Example Tests (Labs)	Common CPT Codes	Common ICD Codes	<u>Ref</u>
<u>Comprehensive</u> <u>Kidney Disease</u> <u>Panels</u>		81401, 81402, 81403, 81404, 81405, 81406, 81407, 81408, 81479	N10-N19,	
	KidneySeq Version 5 Comprehensive Testing (Iowa Institute of Human Genetics)			
	RenalZoom (DNA Diagnostic Laboratory - Johns Hopkins Hospital)			
APOL1-Mediated	Kidney Disease			
<u>Targeted Variant</u> <u>Analysis</u>	Apolipoprotein L1 (APOL1) Renal Risk Variant Genotyping (Quest Diagnostics)	0355U	N00-N08, N10-N19	9
	APOL1 Genotype, Varies (Mayo Clinic Laboratories)	81479		
Donor-Derived Co	ell Free DNA for Kidney Transplant R	ejection		
<u>Donor-Derived</u> <u>Cell Free DNA for</u> <u>Kidney</u> <u>Transplant</u> <u>Rejection</u>	Allosure Kidney (CareDx, Inc.)	81479	T86.11, T86.12,	7
	Prospera Kidney (Natera)		Z94.0	
	Viracor TRAC Kidney dd-cfDNA (Viracor Eurofins)	0118U		
Other Covered K	idney Disorders			
<u>Other Covered</u> <u>Kidney Disorders</u>	See list below	81400, 81401, 81402, 81403, 81404, 81405, 81406, 81407, 81408, 0268U		4, 5, 6

# **OTHER RELATED POLICIES**

This policy document provides coverage criteria for hereditary kidney disorders. Please refer to:

<u>Genetic Testing: Multisystem Inherited Disorders, Intellectual Disability, and</u>
<u>Developmental Delay MP9587</u> for coverage criteria related to genetic disorders
that affect multiple organ systems



- Genetic Testing: Hereditary Cancer Susceptibility MP9596 for coverage criteria related to von Hippel Lindau (VHL) syndrome and other hereditary cancer syndromes.
- Genetic Testing: General Approach to Genetic and Molecular Testing MP9610 for coverage criteria related to genetic testing for kidney disease that is not specifically discussed in this or another non-general policy.

# **COVERAGE CRITERIA**

## POLYCYSTIC KIDNEY DISEASE

### **Targeted Variant Analysis**

- PKD1, PKD2, GANAB, or DNAJB11 targeted variant analysis (81403) to Ι. establish a diagnosis of autosomal dominant polycystic kidney disease is considered medically necessary when:
  - A. The member has a close relative with a known pathogenic or likely pathogenic variant in PKD1, PKD2, GANAB, or DNAJB11.
- Π. *PKHD1* targeted variant analysis (81403) to establish a diagnosis of autosomal recessive polycystic kidney disease is considered medically necessary when:
  - A. The member has a biological full sibling with known biallelic pathogenic or likely pathogenic variants in *PKHD1*.
- 111. PKD1, PKD2, GANAB, DNAJB11, or PKHD1 targeted variant analysis (81403) to establish a diagnosis of autosomal dominant or autosomal recessive polycystic kidney disease is considered investigational for all other indications.

### Single Gene or Multigene Panel

- PKD1 (81407, 81479), PKD2 (81406, 81479), GANAB (81479), DNAJB11 Ι. (81479), PKHD1 (81408, 81479) sequencing and/or deletion/duplication analysis or multigene panel analysis (81404, 81405, 81406, 81407, 81408, 81479) to confirm or establish a diagnosis of polycystic kidney disease is considered medically necessary when:
  - A. The member has any of the following clinical features of polycystic kidney disease.
    - 1. Multiple bilateral renal cysts, OR
    - 2. Cysts in organs other than the kidneys (especially the liver, seminal vesicles, pancreas, and arachnoid membrane), OR



- 3. Hypertension in an individual younger than age 35, OR
- 4. Intracranial aneurysm, **OR**
- 5. Bilaterally enlarged and diffusely echogenic kidneys, OR
- 6. Poor corticomedullary differentiation, OR
- 7. Hepatobiliary abnormalities with progressive portal hypertension, **OR**
- 8. Congenital hepatic fibrosis (CHF) with portal hypertension.
- II. PKD1 (81407, 81479), PKD2 (81406, 81479), GANAB (81479), DNAJB11 (81479), PKHD1 (81408, 81479) sequencing and/or deletion/duplication analysis or multigene panel analysis (81404, 81405, 81406, 81407, 81408, 81479) to confirm or establish a diagnosis of polycystic kidney disease is considered investigational for all other indications.

back to top

### COMPREHENSIVE KIDNEY DISEASE PANELS

- I. Genetic testing for kidney disease via a comprehensive kidney disease panel (81401, 81402, 81403, 81404, 81405, 81406, 81407, 81408, 81479) is considered **medically necessary** when:
  - A. The member has chronic kidney disease with an undetermined cause after undergoing standard-of-care workup studies (examples: history and physical examination, biochemical testing, renal imaging, or renal biopsy), **AND**
  - B. The member meets at least one of the following:
    - 1. Onset of chronic kidney disease under 40 years of age, OR
    - 2. One or more <u>first- or second-degree relatives</u> with chronic kidney disease, **OR**
    - 3. Consanguineous family history, OR
    - 4. Cystic renal disease, OR
    - 5. Congenital nephropathy.
- II. Genetic testing for kidney disease via a comprehensive kidney disease panel (81401, 81402, 81403, 81404, 81405, 81406, 81407, 81408, 81479) is considered **investigational** for all other indications.



back to top

### APOL1-MEDIATED KIDNEY DISEASE

### **Targeted Variant Analysis**

- I. Targeted variant analysis for the *APOL1* high-risk genotype (i.e., G1/G1, G1/G2, or G2/G2) (0355U, 81479) is considered **medically necessary** when:
  - A. The member has kidney disease, AND
    - 1. The member is of African ancestry, **OR**
    - 2. The member has a family member with a confirmed *APOL1* high-risk genotype (i.e., G1/G1, G1/G2, or G2/G2)
- II. Targeted variant analysis for the *APOL1* high-risk genotype (i.e., G1/G1, G1/G2, or G2/G2) (0355U, 81479) is considered **investigational** for all other indications.

back to top

### DONOR-DERIVED CELL-FREE DNA FOR KIDNEY TRANSPLANT REJECTION

- I. The use of peripheral blood measurement of donor-derived cell-free DNA in the management of patients after renal transplantation (81479, 0118U) is considered **investigational** for all indications, including but not limited to:
  - A. Detection of acute renal transplant rejection
  - B. Detection of renal transplant graft dysfunction

back to top

### OTHER COVERED KIDNEY DISORDERS

The following is a list of conditions that have a known genetic association. Due to their relative rareness, it may be appropriate to cover these genetic tests to establish or confirm a diagnosis.

I. Genetic testing to establish or confirm one of the following genetic kidney disorders to guide management is considered **medically necessary** when the member demonstrates clinical features\* consistent with the disorder (the list is not meant to be comprehensive, see II below):



- A. Alport Syndrome
- B. C3 Glomerulopathy
- C. Congenital nephrotic syndrome
- D. Cystinosis
- E. Cystinuria
- F. Fabry Disease
- G. Genetic (familial) atypical hemolytic-uremic syndrome (aHUS)
- H. Primary Hyperoxaluria
- II. Genetic testing to establish or confirm the diagnosis of all other kidney disorders not specifically discussed within this or another medical policy will be evaluated by the criteria outlined in *General Approach to Genetic and Molecular Testing* (see policy for coverage criteria).

\*Clinical features for a specific disorder may be outlined in resources such as <u>GeneReviews</u>, <u>OMIM</u>, <u>National Library of Medicine</u>, <u>Genetics Home Reference</u>, or other scholarly source.

back to top

# **NOTES AND DEFINITIONS**

- 1. Close relatives include first, second, and third degree <u>blood</u> relatives on the same side of the family:
  - a. First-degree relatives are parents, siblings, and children
  - b. **Second-degree relatives** are grandparents, aunts, uncles, nieces, nephews, grandchildren, and half siblings
  - c. **Third-degree relatives** are great grandparents, great aunts, great uncles, great grandchildren, and first cousins
- 2. Full siblings are individuals who share the same biological parents.

# **BACKGROUND AND RATIONALE**

#### Polycystic Kidney Disease - Targeted Variant Analysis

Genetic Support Foundation

The Genetic Support Foundation's Genetics 101 information on genetic testing says the following about testing for familial pathogenic variants:

Genetic testing for someone who may be at risk for an inherited disease is always easier if we know the specific genetic cause. Oftentimes, the best way to find the



genetic cause is to start by testing someone in the family who is known or strongly suspected to have the disease. If their testing is positive, then we can say that we have found the familial pathogenic (harmful) variant. We can use this as a marker to test other members of the family to see who is also at risk.

#### Polycystic Kidney Disease - Single Gene or Multigene Panel

*GeneReviews: Polycystic Kidney Disease, Autosomal Dominant and Polycystic Kidney Disease, Autosomal Recessive* 

GeneReviews is an expert-authored review of current literature on a genetic disease, and goes through a rigorous editing and peer review process before being published online. The recommended polycystic kidney disease testing for autosomal dominant polycystic kidney disease (ADPKD) and autosomal recessive polycystic kidney disease (ARPKD) is as follows:

"ADPKD should be suspected in individuals with the following:

- Multiple bilateral renal cysts and the absence of manifestations suggestive of a different renal cystic disease
- Cysts in other organs, especially the liver, but also seminal vesicles, pancreas, and arachnoid membrane...
- Hypertension in an individual younger than age 35 years
- An intracranial aneurysm..."

"Autosomal recessive polycystic kidney disease (ARPKD) should be suspected in individuals with bilaterally enlarged, diffusely echogenic kidneys...[and] one or more of the following:...Clinical/laboratory signs of congenital hepatic fibrosis (CHF) that leads to portal hypertension..."

"The renal diagnostic criteria for ARPKD detected by ultrasonography are:

- Increased renal size (in relation to normative size based on age and size of the affected individual);
- Increased echogenicity;
- Poor corticomedullary differentiation"

"[In] Childhood and young adulthood...The hepatobiliary abnormalities with progressive portal hypertension are often the prominent presenting features."

#### **Comprehensive Kidney Disease Panels**

Hays et al (2020)



"We propose the following approach, based on a review of current literature and our practical experience. This approach assumes individuals have already undergone an initial nephrologic workup, including biochemical and serologic testing, imaging of the kidneys, and renal biopsy if indicated.

...[A]fter a negative or inconclusive initial workup, a patient is considered to have KDUE [kidney disease of unknown etiology] and may then be stratified according to the probability of a genetic disease. We consider higher probability patients as those with the following risk factors: early-onset disease (age <40 years), a positive family history of CKD [chronic kidney disease], consanguinity, extrarenal anomalies, cystic renal disease, or congenital nephropathy". (p. 594)

#### **APOL1-Mediated Kidney Disease**

#### Freedman et al (2021)

A multidisciplinary group of experts and patient advocates performed a systematic review and created consensus-based guidelines in 2021 to guide health care providers in *APOL1*-associated neuropathy. The guidelines recommend the following:

"...*APOL1* testing should be considered in all patients of African ancestry with kidney disease and in any patient with kidney disease and a family member with a confirmed *APOL1* high-risk genotype." (p. 1768)

Regarding the definition of "high-risk phenotype": "Two copies of the *APOL1* variants (G1/G1, G1/G2, G2/G2) are commonly referred to as a 'high-risk' genotype…" (p. 1765)

#### **Donor-Derived Cell-Free DNA for Kidney Transplant Rejection**

Knight et al (2019)

A publication in the journal Transplantation entitled "Donor-specific Cell-free DNA as a Biomarker in Solid Organ Transplantation. A Systematic Review" stated the following:

In summary, donor-derived cfDNA shows promise as a biomarker for the detection of acute transplant graft injury. It has potential to reduce the need for protocol biopsy surveillance, allowing for a more targeted diagnostic approach. Detection of injury occurs before clinical manifestation, meaning that there is a window for earlier detection and treatment of AR [acute rejection] and other causes of graft injury with the potential to improve outcomes. It may also facilitate the detection of under immunosuppression and find use as a tool for monitoring during immunosuppression minimization. Further studies are required to validate the thresholds for further investigation and intervention, determine the optimum frequency for monitoring, and to identify whether prospective monitoring using



dd-cfDNA can indeed improve transplant outcomes compared to current practice. (p. 280)

back to top

## REFERENCES

- Harris PC, Torres VE. Polycystic Kidney Disease, Autosomal Dominant. 2002 Jan 10 [Updated 2022 Sep 29]. In: Adam MP, Ardinger HH, Pagon RA, et al., editors. GeneReviews [Internet]. Seattle (WA): University of Washington, Seattle; 1993-2023. Available from: <u>https://www.ncbi.nlm.nih.gov/books/NBK1246/</u>
- Sweeney WE, Avner ED. Polycystic Kidney Disease, Autosomal Recessive. 2001 Jul 19 [Updated 2019 Feb 14]. In: Adam MP, Ardinger HH, Pagon RA, et al., editors. GeneReviews [Internet]. Seattle (WA): University of Washington, Seattle; 1993-2023. Available from: https://www.ncbi.nlm.nih.gov/books/NBK1326/
- Hays T, Groopman EE, Gharavi AG. Genetic testing for kidney disease of unknown etiology. Kidney Int. 2020;98(3):590-600. doi:10.1016/j.kint.2020.03.031
- Adam MP, Ardinger HH, Pagon RA, et al., editors. GeneReviews [Internet]. Seattle (WA): University of Washington, Seattle; 1993-2023. Available from: <u>https://www.ncbi.nlm.nih.gov/books/NBK1116/</u>
- Online Mendelian Inheritance in Man, OMIM. McKusick-Nathans Institute of Genetic Medicine, Johns Hopkins University (Baltimore, MD). World Wide Web URL: <u>https://omim.org/</u>
- 6. MedlinePlus [Internet]. Bethesda (MD): National Library of Medicine (US). Available from: <u>https://medlineplus.gov/genetics/</u>.
- Knight SR, Thorne A, Lo Faro ML. Donor-specific cell-free DNA as a biomarker in solid organ transplantation. A systematic review. *Transplantation*. 2019;103(2):273-283.
- Genetic Support Foundation. Genetics 101 Genetic Testing: Familial Pathogenic Variant. Accessed 10/4/2022. <u>https://geneticsupportfoundation.org/genetics-101/#</u>
- 9. Freedman BI, Burke W, Divers J, et al. Diagnosis, education, and care of patients with APOL1-associated nephropathy: a Delphi consensus and systematic review. JASN. 2021;32(7):1765-1778.

back to top



### **Committee/Source**

### Date(s)

Document Created:	Medical Policy Committee/Health Services Division	November 16, 2022
Revised:	Medical Policy Committee/Health Services Division Medical Policy Committee/Health Services Division Medical Policy Committee/Health Services Division	February 15, 2023 March 15, 2023 August 16, 2023
Reviewed:	Medical Policy Committee/Health Services Division Medical Policy Committee/Health Services Division Medical Policy Committee/Health Services Division	February 15, 2023 March 15, 2023 August 16, 2023

Published: 01/01/2024 Effective: 01/01/2024